

## **EMERGENCY CONTRACEPTION OPTIONS 250-592-3479**

### islandsexualhealth.org

#### What is Emergency Contraception?

Emergency Contraception (EC) is contraception that is used to reduce the risk of unintentional pregnancy after either unprotected vaginal intercourse or a known/suspected birth control failure such as a broken/slipped condom, missed pill, delayed insertion of ring, detached patch, late injection, or no use of birth control. The effectiveness of EC will vary depending on the type of method used and timing of use. The Copper IUD is the most effective EC method to reduce risk of unintentional pregnancy and has the longest timeframe (w/i 7days) to utilize while Emergency Contraceptive Pills (ECP) vary in effectiveness and time frame depending on type.

#### There are 2 EC options available:

- Copper IUD
- Emergency Contraceptive Pill (ECP) Ullipristal Acetate or Progestin Only (see 2nd page)

Progestin Pill (Plan B, Contingency One...)

Ullipristal Acetate (Ella)

**Copper IUD** 

Moderatively Effective ~50%

Moderatively Effective ~60%

**Most Effective 99%** 

Type of EC	Effectiveness?	Advantages?	How do I use it ?	User Weight Limit?	Where can I Get it?
Copper IUD	99% if inserted within 7 days after vaginal intercourse.	User can choose to leave the IUD in place as a very effective (99%) ongoing birth control method for 5 to 10 years depending on the IUD model.	A small T shaped device containing copper is placed into the uterus by a health care provider.	None	<ul> <li>By Prescription and Insertion through:</li> <li>Island Sexual Health 250-592-3479</li> <li>VI Women's Clinic 250-480-7338</li> <li>Women's Vita Clinic (Nanaimo) 250-591-9812</li> </ul>
	Time Frame for Use?  Within 7 days after vaginal intercourse.	Disadvantages?  Must be prescribed/ inserted by a health care provider  Cost  May increase cramping and flow for longer term users	Copper creates an inflammatory reaction in the uterus which prevents fertilization and/or implantation.	\$100-150 for device No cost for insertion if user has MSP	<ul> <li>emergencyiud.com (Listing of clinics providing emergency iud insertions in Western Canada)</li> <li>raice.ca (Canada wide listing of rapid access IUD clinics)</li> <li>Turn page over for info on Emergency Contraceptive Pills</li> </ul>



# **Emergency Contraception Options - Emergency Contraceptive Pill** islandsexualhealth.org

ECP only decreases the risk of pregnancy occurring; it does not end an established pregnancy. It is <u>not</u> an abortion pill.

Туре	Effectiveness?	Advantages?	How do I use it ?	User	Where Can I Get it?	
of EC				Weight Limit?		
Ullipristal Acetate sold as (Ella®)	~64% if taken within 120 hours after single act of unprotected vaginal intercourse.  Time Frame for Use?	Slightly more effective than progestin pill.  Equally effective up to 120 hrs following unprotected intercourse.  Disadvantages?  Ella interferes with progesterone	1 Pill taken by mouth  How does it work?	May be <i>more</i> effective in users who weigh more than 165lbs or have a higher BMI.	Requires a prescription from a Dr. or Nurse Practitioner.  Available through prescription at:  Island Sexual Health 3960 Quadra St. 250-592-3479	
R	As soon as possible.  Must be used within 120 hrs (5 days).	so it's not the best choice for clients who are using it with pill, patch, ring or shot.  Ongoing hormonal contraception cannot be started for 5 days after using Ullipristal. Back up with a barrier method for 14 days.  Does not work if ovulation has already occurred.	ECP works by delaying ovulation so there is an increased risk of pregnancy if user has unprotected sex after taking ECP without using a back up method before their next period.  *If breastfeeding, speak with a Dr./Nurse/pharmacist prior to Ella use.*	\$25-40/per dose	<ul> <li>Women's Vita Clinic         (Nanaimo) 250-591-9812</li> <li>GPs/Walk in clinics (take in prescription to pharmacy to be filled. Call pharmacy ahead to confirm availability)</li> </ul>	
Progestin Pill  (Plan B®, Contingency one®, Next Choice®, Backup Plan OneStep®, Norlevo®)	~50%  Most effective if taken up to 72 hrs after unprotected vaginal intercourse. Effectiveness decreases over time but may be taken up to 120 hrs following.  Time Frame for Use?  As soon as possible. Ideally within 72 hrs. Must be used within 120 hrs (5 days)	No prescription necessary.  May be purchased ahead of time to have available.  Ongoing hormonal contraception can be started immediately after taking ECP. Back up with a barrier method for the first 7 days.  Disadvantages?  Effectiveness of progestin only pill declines as the time increases between intercourse and use.  Does not work if ovulation has already occurred.	1 or 2 Pills taken by mouth  How does it work?  ECP works by delaying ovulation so there is an increased risk of pregnancy if user has unprotected sex after taking ECP without using a back up method before their next period. *Safe to use if breastfeeding.*	May be less effective in users who weigh more than 165lbs. Cost? \$17-\$45/per dose	Prescription not needed.  Available to purchase by/for anyone:  Island Sexual Health 3960 Quadra St. 250-592-3479  Youth Clinics  Available at Most Pharmacies  planb.ca (listing of clinics across Canada)	